

Clarithro

clarithromycin

Consumer Medicine Information

What is in this leaflet

This leaflet answers some common questions about CLARITHRO.

It does not contain all of the available information. It does not take the place of talking to your doctor or pharmacist.

All medicines have benefits and risks. Your doctor has weighed the risks of you using CLARITHRO against the benefits they expect it will have for you.

If you have any concerns about taking this medicine, talk to your doctor or pharmacist.

Keep this leaflet with your medicine. You may need to read it again.

What CLARITHRO is used for

CLARITHRO is used to treat certain bacterial infections, including the following:

- respiratory tract infections
- skin infections
- peptic ulcer.

CLARITHRO is also used to prevent a specific bacterial infection associated with HIV infection.

Your doctor, however, may have prescribed CLARITHRO for another purpose.

Ask your doctor if you have any questions about why CLARITHRO has been prescribed for you.

CLARITHRO is an antibiotic that belongs to the group of medicines called macrolides. These medicines

work by killing or stopping the growth of bacteria which cause infections.

CLARITHRO will not work against infections caused by viruses, such as colds or flu.

CLARITHRO is available only with a doctor's prescription.

CLARITHRO to treat peptic ulcer

Peptic ulcers are associated with an infection in the intestine and stomach caused by the bacteria *Helicobacter pylori* (*H. pylori*). Nearly all patients with peptic ulcers are infected with *H. pylori*.

The *H. pylori* infection can be treated with a combination of CLARITHRO (clarithromycin), another antibiotic (amoxicillin) and another medicine called omeprazole (used to control the acidity of the stomach).

However, the best combination of tablets to treat *H. pylori* infection is yet to be determined. Your doctor will determine the best combination for you.

If your symptoms return, consult your doctor.

It is possible that CLARITHRO may no longer be effective in killing the *H. pylori* infection and a different antibiotic may be needed.

Before you take CLARITHRO

When you must not take it

Do not take CLARITHRO if you are allergic to medicines containing clarithromycin or any

other macrolide antibiotic (such as erythromycin, roxithromycin and azithromycin) or any of the ingredients listed at the end of this leaflet.

Some of the symptoms of an allergic reaction may include rash, itching or hives, swelling of the face, lips, tongue or other parts of the body, shortness of breath, wheezing or difficulty in breathing.

Do not take CLARITHRO if you are taking these medicines:

- astemizole or terfenadine (commonly used to treat allergy symptoms - these medicines may be available without a prescription)
- cisapride (Prepulsid), a medicine used to treat stomach disorders
- domperidone (used for nausea)
- pimozone (Orap), a medicine used to treat psychotic disorders
- ergotamine (Cafergot) or dihydroergotamine (Dihydergot), medicines for migraines
- lovastatin or simvastatin (used to treat high cholesterol)
- ticagrelor or ranolazine (used to prevent blood clotting)
- colchicine (used to treat gout)
- midazolam tablets or syrup (used to produce drowsiness before certain procedures or surgery)

Do not take CLARITHRO if you have:

- severe liver problems
- poor kidney function.

Do not take CLARITHRO if the expiry date (Exp.) printed on the pack has passed.

Do not take CLARITHRO if the packaging shows signs of

and take your next dose when you are meant to.

Otherwise, take the missed dose as soon as you remember, and then go back to taking your tablets as you would normally. Do not take a double dose to make up for the dose you missed.

If you are not sure what to do, ask your doctor or pharmacist.

How long to take it for

Keep taking CLARITHRO until you finish the pack or for as long as your doctor tells you.

If you are being treated for an infection, CLARITHRO is usually taken for one or two weeks.

Do not stop taking CLARITHRO, even if you feel better after a few days, unless advised by your doctor.

Your infection may not clear completely if you stop taking your medicine too soon.

Check with your doctor if you are not sure how long you should be taking CLARITHRO.

If you take too much CLARITHRO (overdose)

Immediately telephone your doctor, or the Poisons Information Centre (telephone 13 11 26), or go to Accident and Emergency at the nearest hospital, if you think you or anyone else may have taken too much CLARITHRO. Do this even if there are no signs of discomfort or poisoning.

You may need urgent medical attention.

If you take too much CLARITHRO, you may develop severe gastrointestinal symptoms, liver problems, or allergic reactions.

While you are taking CLARITHRO

Things you must do

If you are taking CLARITHRO for an infection and your symptoms do not improve within a few days, or if they become worse, tell your doctor.

Tell your doctor if you become pregnant while taking CLARITHRO.

If you get severe diarrhoea, tell your doctor or pharmacist immediately.

Do this even if it occurs several weeks after stopping CLARITHRO. Diarrhoea may mean that you have a serious condition affecting your bowel. You may need urgent medical care. Do not take any medicine to stop your diarrhoea without first checking with your doctor.

If you have to have any urine tests, tell your doctor you are taking CLARITHRO as it may affect the results of some laboratory tests.

Before starting any new medicine, tell your doctor or pharmacist that you are taking CLARITHRO.

Tell all the doctors, dentists and pharmacists who are treating you that you are taking CLARITHRO.

Things you must not do

Do not use CLARITHRO to treat any other conditions unless your doctor tells you to.

Do not give CLARITHRO to anyone else, even if they have the same condition as you.

Side effects

Tell your doctor or pharmacist as soon as possible if you do not feel well while you are taking CLARITHRO.

CLARITHRO treats infections in most people, but it may have unwanted side effects in some people.

All medicines can have side effects. Sometimes they are serious, most of the time they are not. You may need medical treatment if you get some of the side effects.

Do not be alarmed by this list of possible side effects.

You may not experience any of them.

Ask your doctor or pharmacist to answer any questions you may have.

Tell your doctor if you notice any of the following and they worry you:

- stomach cramps and pains
- nausea, vomiting and severe diarrhoea
- oral thrush or vaginal thrush
- change in taste sensation
- headache
- asthma, shortness of breath
- muscle weakness
- conjunctivitis.

Tell your doctor immediately if you notice any of the following:

- yellowing of the eyes or skin (jaundice)
- feeling generally unwell and having poor appetite
- hearing disturbances
- chest pain
- dizziness, confusion, hallucinations, convulsions
- fainting, irregular heartbeat
- any type of skin rash, itching, hives
- severe diarrhoea, especially if bloody
- severe upper stomach pain, with nausea and vomiting (pancreatitis).

These are serious side effects. You may need urgent medical treatment.

If any of the following happen, stop taking CLARITHRO and tell your doctor immediately, or

go to Accident and Emergency at the nearest hospital:

- symptoms of an allergic reaction such as swelling of the face, lips, mouth or throat, which may cause difficulty in swallowing or breathing, shortness of breath or sudden collapse.

After using CLARITHRO

Tell your doctor immediately if you notice any of the following side effects, even if they occur several weeks after stopping treatment with CLARITHRO:

- severe stomach or abdominal cramps
- watery and severe diarrhoea, which may also be bloody
- fever, in combination with one or both of the above.

These are rare but serious side effects. You may have a serious condition affecting your bowel and you may need urgent medical care.

Do not take any diarrhoea medicine without first checking with your doctor.

Other side effects not listed above may also occur in some patients. Tell your doctor if you notice anything that is making you feel unwell.

After using CLARITHRO

Storage

Keep CLARITHRO where children cannot reach it.

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

Keep CLARITHRO in a cool dry place, protected from light, where the temperature stays below 25°C.

Do not store CLARITHRO or any other medicine in the bathroom or near a sink.

Do not leave CLARITHRO in the car or on window sills.

Heat and dampness can destroy some medicines.

Disposal

If your doctor tells you to stop using CLARITHRO, or your medicine has passed its expiry date, ask your pharmacist what to do with any tablets that are left over.

Product description

What it looks like

CLARITHRO is a yellow, oval film-coated tablet.

Each blister pack contains 14 tablets.

Ingredients

The active ingredient in CLARITHRO is Clarithromycin. Each CLARITHRO 250 tablet contains 250 mg of Clarithromycin.

CLARITHRO tablets also contain:

- microcrystalline cellulose
- croscarmellose sodium
- pregelatinised maize starch
- povidone
- purified talc
- magnesium stearate
- colloidal anhydrous silica
- Opadry II 40L22445 yellow.

Sponsor

Arrow Pharma Pty Ltd
15-17 Chapel Street
Cremorne VIC 3121

Australian registration numbers:
CLARITHRO 250 - AUST R
117197

Date of preparation:
August 2018