

# OXYTOCIN GH INJECTION

(oxytocin) solution for injection

---

## Consumer Medicine Information

---

### What is in this leaflet

---

This leaflet answers some common questions about Oxytocin GH Injection.

It does not contain all the available information.

It does not take the place of talking with your doctor or pharmacist.

The information in this leaflet was last updated on the date listed on the final page. Some more recent information on the medicine may be available.

**You should ensure that you speak to your pharmacist or doctor to obtain the most up to date information on the medicine.**

Those updates may contain important information about the medicine and its use of which you should be aware.

All medicines have risks and benefits. Your doctor has weighed the risks of your treatment using Oxytocin GH Injection against the benefits they expect it will have for you.

**If you have any concerns about taking this medicine, ask your doctor or pharmacist.**

**Keep this leaflet with the medicine.**

You may need to read it again

---

### What OXYTOCIN GH INJECTION is used for

---

Oxytocin GH Injection can be used to bring on (induce) labour. It can also be used during and immediately after delivery to help

the birth and to prevent or treat excessive bleeding.

Oxytocin GH Injection is a man-made chemical that is identical to a natural hormone called oxytocin. It works by stimulating the muscles of the uterus (womb) to produce rhythmic contractions.

Oxytocin GH Injection is not suitable in all situations - for example, if the baby or placenta are in the wrong position or if you have had a previous caesarean section or other surgery involving the uterus. Your doctor can give you more information on the suitability of Oxytocin GH Injection in your particular case.

**Ask your doctor if you have any questions about why Oxytocin GH Injection has been prescribed for you.**

Your doctor may have prescribed it for another purpose.

Oxytocin GH Injection is only available with a doctor's prescription. It is not addictive.

---

### Before OXYTOCIN GH INJECTION is given

---

*When Oxytocin GH Injection should not be used*

**You must not have Oxytocin GH Injection if:**

- your doctor thinks that inducing or enhancing contractions for normal labour and vaginal delivery would be unsuitable for you or your baby.
- there are maternal or foetal reasons for caesarean delivery.

- you have been given medicines called prostaglandins within the past 6 hours.

**You must not have Oxytocin GH Injection if you are allergic to:**

- oxytocin (the active ingredient)
- any of the other ingredients listed at the end of this leaflet.

Symptoms of an allergic reaction may include:

- shortness of breath
- wheezing or difficulty breathing
- swelling of the face, lips, tongue or other parts of the body
- rash, itching or hives on the skin.

*Before you are given it*

**Tell your doctor if you have high blood pressure or any heart or kidney problems.**

Your doctor may want to take extra precautions. For example, the amount of fluid you will be given may need to be reduced if you have a problem with your heart or kidneys.

**Tell your doctor if you are allergic to any other medicines, foods, dyes or preservatives.**

Your doctor will want to know if you are prone to allergies.

**Tell your doctor if you have, or have ever had, any of the following problems:**

- an abnormal electrical signal called "prolongation of the QT interval";
- any other conditions that affect the heart;
- kidney problems.

**Tell your doctor if you are taking any medicines that may affect**

**your heart, or any other medicines, including medicines that you buy without a prescription from a pharmacy, supermarket or health food shop.**

**Tell your doctor if you have been given anaesthetics or medicines called prostaglandins.**

**If you have not told your doctor about any of these things, tell him/her before you have Oxytocin GH Injection.**

---

## How OXYTOCIN GH INJECTION is given

---

To bring on (induce) or maintain labour, Oxytocin GH Injection is given by intravenous infusion (drip). The speed of the infusion is set to maintain a pattern of contractions similar to normal labour. During the infusion, both you and your baby will be closely monitored to prevent complications.

If Oxytocin GH Injection is needed at delivery or to prevent excessive bleeding, it can also be given intramuscularly (into a muscle) or by slow intravenous injection directly into a vein.

---

## Side effects

---

**Tell your doctor or nurse as soon as possible if you do not feel well while you are having Oxytocin GH Injection.**

All medicines can have side effects. Sometimes they are serious, most of the time they are not. You may need medical treatment if you get some of the side effects.

Tell your doctor or nurse immediately if you notice any of the following symptoms:

- rash, itching or hives on the skin;
- swelling of the face, lips, tongue or other parts of the body;

- shortness of breath, wheezing or troubled breathing;
- headache;
- nausea (feeling sick) or vomiting;
- feeling drowsy and lethargic;
- pain in the abdomen that is different from labour pains;
- dizziness, light headedness or faintness;
- flushing of the face;
- chest pain;
- fast, slow or irregular heart beat;
- excessive or continuous contractions;
- abnormal clotting or bleeding.

The above symptoms may be signs of allergy or signs of too much fluid associated with high doses or long infusions.

**Tell your doctor if you notice anything else that is making you feel unwell.**

Other side effects not listed above may happen in some people.

---

## After using OXYTOCIN GH INJECTION

---

### *Storage*

**Store unopened Oxytocin GH Injection ampoules in a refrigerator, where the temperature stays between 2°C-8°C. Protect from light and do not freeze.**

The unopened product may be stored for up to three (3) months at or below 25°C. Once removed from the refrigerator the product must be discarded if not used. It must not be returned to the refrigerator.

Oxytocin GH Injection will be kept in a locked cupboard in the hospital pharmacy or operating theatre.

Oxytocin GH Injection should not be used after the date (month and year) printed after "EXP". The anaesthetist will inspect Oxytocin GH Injection before use to

determine that it is still within its use by date.

### *Disposal*

The hospital staff looking after you will dispose of any remaining Oxytocin GH Injection appropriately.

---

## Product description

---

### *What it looks like*

Oxytocin GH Injection is a clear, colourless solution.

Oxytocin GH Injection is available 1 mL glass ampoules containing either 8.3 microgram (5 IU) or 16.7 micrograms (10 IU) of oxytocin.

A carton of Oxytocin GH Injection contains 1, 5 or 10 ampoules. Not all pack sizes may be marketed.

### *Ingredients*

#### **Active ingredient:**

- oxytocin

#### **Other ingredients:**

- glacial acetic acid
- sodium acetate
- sodium chloride
- water for injections

### *Australian Registration Numbers*

5 IU/1 mL - AUST R 207985

10 IU/1 mL - AUST R 207986

### *Distributor*

Generic Health Pty Ltd  
Suite 2, Level 2  
19-23 Prospect Street  
Box Hill, VIC, 3128  
Australia

Email:

ghinfo@generichealth.com.au

Phone: +61 3 9809 7900

Website:

www.generichealth.com.au

This leaflet was prepared in August 2018.