

Lidocaine (lignocaine) hydrochloride Injection BP

Consumer Medicine Information

What is in this leaflet

This leaflet answers some common questions about Lidocaine (lignocaine) hydrochloride Injection. It does not contain all the available information. It does not take the place of talking to your doctor or pharmacist.

All medicines have benefits and risks. Your doctor has weighed the risks of you taking Lidocaine (lignocaine) hydrochloride Injection against the benefits this medicine is expected to have for you.

This medicine is likely to be used while you are at the clinic or in hospital. If possible, please read this leaflet carefully before this medicine is given to you. In some cases this leaflet may be given to you after the medicine has been used.

If you have any concerns about taking this medicine, ask your doctor or pharmacist.

Keep this leaflet.

You may need to read it again.

What Lidocaine (lignocaine) hydrochloride Injection is used for

Lidocaine (lignocaine) hydrochloride Injection belongs to two groups of medicines known as local anaesthetics and antiarrhythmic drugs. Local anaesthetics stop pain and feeling in the area around where it is injected;

and antiarrhythmic drugs work by restoring irregular and/or rapid heart beats to normal.

Lidocaine (lignocaine) hydrochloride Injection may be used for the management of other conditions that are not mentioned above. Your doctor will be able to tell you about the specific condition for which you have been prescribed Lidocaine (lignocaine) hydrochloride Injection.

This medicine is available only with a doctor's prescription.

Before you are given Lidocaine (lignocaine) hydrochloride Injection

When you must not be given it

Do not use Lidocaine (lignocaine) hydrochloride Injection if:

- you have an allergy to lignocaine or any of the ingredients listed at the end of this leaflet. Symptoms of an allergic reaction may include shortness of breath, wheezing or difficulty breathing, swelling of the face, lips, tongue or other parts of the body, rash, itching or hives on the skin.
- you have an infection or swelling where the injection will be given
- you have blood infections or disorders
- you are in severe shock
- you have low blood pressure

- you have diseases of the brain and spinal cord
- you have sudden frequent episodes of loss of consciousness
- you have myasthenia gravis, a severe muscle weakness
- you have certain heart problems

If you are not sure whether any of these apply to you, check with your doctor.

Before you are given it

Tell your doctor if:

1. you have any allergies to:

- any other medicine
- any other substances, such as foods, preservatives or dyes

2. you are pregnant or intend to become pregnant

3. you are breast-feeding or plan to breast feed

4. you have or have had any medical conditions, especially the following:

- liver and/or kidney problems
- epilepsy
- diseases of the brain and spinal cord
- heart problems
- severe fever
- problems with your blood pressure or circulation
- slow heart beat
- low potassium level
- breathing difficulties
- low oxygen in blood
- problems with the clotting of your blood

Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from your pharmacy, supermarket or health food shop.

Some medicines and lignocaine may interfere with each other.

These include:

- anti-arrhythmics, medicines to control irregular heart beats, e.g. amiodarone
- heart or blood pressure tablets, e.g. propranolol, metoprolol, atenolol
- cimetidine, a medicine used to treat reflux and ulcers
- medicines to control fits or epilepsy, e.g. phenytoin, carbamazepine, primidone, phenobarbitone. sodium valproate
- warfarin, a medicine used to stop blood clots
- aspirin, salicylates or other non-steroidal anti-inflammatory drugs (NSAIDS), medicines to relieve pain and inflammation
- some anaesthetics
- some medicines that relax the skeletal muscles, e.g. suxamethonium
- nitrates/nitrites e.g. nitric oxide, nitroglycerin, sodium nitroprusside, nitrous oxide
- antineoplastic agents e.g. cyclophosphamide monohydrate, flutamide, hydroxyurea, ifosfamide, rasburicase
- some antibiotics
- antimalarials e.g. chloroquin phosphate, primaquine
- other drugs such as paracetamol, metoclopramide, quinine, sulfasalazine

Your doctor will have more information on medicines to be careful with or avoid while you are being given Lidocaine (lignocaine) hydrochloride Injection. Your doctor will advise you about

continuing to take other medicines while you are receiving Lidocaine (lignocaine) hydrochloride Injection.

How Lidocaine (lignocaine) hydrochloride Injection is given

Lidocaine (lignocaine) hydrochloride Injection is given by injection into the skin, directly into the blood stream or into an organ. It must only be given by a doctor or nurse.

Your doctor will decide what dose and how long you will receive Lidocaine (lignocaine) hydrochloride Injection.

If you are given too much (overdose)

This rarely happens as Lidocaine (lignocaine) hydrochloride Injection is administered under the care of a highly trained doctor.

However, if you are given too much lignocaine, you may experience some of the effects listed under "Side Effects" below.

Your doctor has information on how to recognise and treat an overdose. Ask your doctor if you have any concerns.

While you are being given Lidocaine (lignocaine) hydrochloride Injection

Things to be careful of

Be careful driving or operating machinery after you have been given Lidocaine (lignocaine) hydrochloride Injection. You may be drowsy or your reflexes may be slow.

Do not drink alcohol while you are being given Lidocaine (lignocaine) hydrochloride Injection. If you drink alcohol while you are being given Lidocaine (lignocaine) hydrochloride Injection, your blood pressure may drop making you feel dizzy and faint.

Side effects

Tell your doctor, nurse or pharmacist as soon as possible if you do not feel well while you are being given lignocaine.

Like other medicines, lignocaine can cause some side effects. If they occur, most are likely to be minor or temporary. However, some may be serious and need medical attention.

Ask your doctor or pharmacist to answer any questions that you may have.

Do not be alarmed by this list of possible side effects. You may not experience any of them.

Tell your doctor or nurse if you notice any of the following:

- weakness, confusion, nervousness, agitation, drowsiness, disorientation or unconsciousness
- nausea, vomiting, difficulty in swallowing
- headache, dizziness or light-headedness especially if you get up suddenly from a lying or sitting position
- ringing in the ears
- slurred speech, sensations of heat and cold
- numbness, tremors, twitching
- slow heart beat, low blood pressure
- breathing difficulties
- blurred or double vision

These are the mild side effects of Lidocaine (lignocaine) hydrochloride Injection.

Serious side effects of Lidocaine (lignocaine) hydrochloride

Injection include:

- fits or convulsions
- skin rash, hives or itching
- unconsciousness
- breathing problems
- low blood pressure
- slow heart beat
- collapse
- pale, grey, or blue coloured skin (cyanosis)
- rapid heart rate

You may need urgent medical attention if you get these side effects.

Other side effects not listed above may also occur in some patients. If you notice any other effects, check with your doctor. Some side effects may only be seen by your doctor.

Storage

Lidocaine (lignocaine) hydrochloride Injection will be stored in the pharmacy or on the ward. The injection is kept in a cool dry place, where the temperature stays below 25 degrees C.

Product description

What it looks like

Lidocaine (lignocaine) hydrochloride Injection is a clear, colourless solution in a polyethylene plastic ampoule.

Lidocaine (lignocaine) hydrochloride Injection can be identified by an Australian Registration Number, which is found on the packaging: AUST R 49296, AUST R 49297, AUST R 49293 and AUST R 49295.

Ingredients

Lidocaine (lignocaine) hydrochloride Injection contains Lignocaine Hydrochloride 1% or 2% as the active ingredient and Sodium Chloride and Water for Injections as the excipients. It does not contain a preservative.

Manufacturer

Pfizer (Perth) Pty Limited
ACN 32 051 824 956
15 Brodie Hall Drive
Bentley WA 6102 Australia

Sponsor and Distributor in Australia:

Pfizer Australia Pty Ltd
Level 17, 151 Clarence Street,
Sydney NSW 2000
Toll Free Number: 1800 675 229
www.pfizer.com.au

This Consumer Medicine Information was written in April 2001.

**Date of most recent amendment:
19 November 2019**