

# LERCADIP

Lercanidipine hydrochloride

## Consumer Medicine Information

### What is in this leaflet

This leaflet answers some of the common questions about Lercadip. It does not contain all the available information. It does not replace seeking advice from your doctor or pharmacist.

#### **Please read this leaflet before you start taking Lercadip.**

If you are helping someone else take Lercadip, please read this leaflet before you give the first dose.

All medicines have risks and benefits. Your doctor has weighed the risks of you taking Lercadip against the benefits this medicine is expected to have for you.

#### **If you have any concerns about taking this medicine, ask your doctor or pharmacist.**

#### **Keep this leaflet with the medicine.**

You may need to read it again.

### What Lercadip is used for

The name of your medicine is Lercadip, which is also known as lercanidipine. Lercadip belongs to a group of medicines called calcium channel blockers (of the dihydropyridine group).

Lercadip lowers high blood pressure, which doctors call hypertension. It works by relaxing some of the blood vessels in the body and reducing resistance to the flow of blood through the blood vessels.

Everyone has blood pressure. This pressure helps get your blood all around your body. Your blood pressure may be different at different times of the day, depending on how busy or worried you are. You have hypertension (high blood pressure) when your blood pressure stays higher than is needed, even when you are calm and relaxed.

There are usually no symptoms of hypertension. The only way of knowing that you have hypertension is to have your blood pressure checked on a regular basis. If high blood pressure is not treated it can lead to serious health problems. You may feel fine and have no symptoms, but eventually hypertension can cause stroke, heart disease and kidney failure. Lercadip helps lower your blood pressure.

Lercadip cannot be obtained without a doctor's prescription.

There is no evidence that Lercadip is addictive.

### Before you take Lercadip

#### *When you must not take it*

#### **Do not take Lercadip if you have an allergy to:**

- lercanidipine or any of the ingredients listed at the end of this leaflet
- any drugs closely related to lercanidipine (such as amlodipine, felodipine or nifedipine)

#### **Do not take Lercadip if you:**

- have severe liver or kidney disease
- are also taking another medicine called ciclosporin

Ask your doctor or pharmacist if you are not sure.

#### **Do not take Lercadip if you are pregnant or intend to become pregnant.**

Like most calcium channel blockers, Lercadip is not recommended for use during pregnancy.

#### **Tell your doctor immediately if you become pregnant while you are taking Lercadip.**

#### **Do not take Lercadip if you are breast-feeding or plan to breast-feed.**

Like most calcium channel blockers Lercadip is not recommended while you are breast-feeding.

#### **Do not take Lercadip if the packaging is torn or shows signs of tampering.**

Do not take Lercadip if the tablets show visible sign of deterioration (for example, are broken or discoloured).

#### **Do not take Lercadip if the expiry date (EXP) printed on the pack has passed.**

If you take this medicine after the expiry date has passed, it may not work (as well).

#### **If you are not sure whether you should start taking Lercadip, talk to your doctor.**

#### **Do not give Lercadip to a child under the age of 18 years.**

Lercadip is not recommended for use in children.

#### *Before you start to take it*

#### **You must tell your doctor:**

1. **if you are allergic to any other medicine or any foods, dyes or preservatives.**
2. **if you are taking other drugs for high blood pressure, such as beta-blockers, diuretics, ACE-inhibitors or angiotensin II receptor antagonists.**
3. **if you have or have ever had any other health problems/medical conditions, including:**
  - liver or kidney disease or are on dialysis
  - certain other heart conditions such as: uncontrolled heart failure, an obstruction to flow of blood from the heart, unstable angina (chest pain or tightness at rest or progressively increasing) or you have had a heart attack one month ago or less and/or if you require a pacemaker

#### **If you have not told your doctor about any of the above or if you are not sure, tell them before you take Lercadip.**

#### *Taking other medicines*

**Tell your doctor if you are taking any other medicines, including medicines that you buy without a prescription from your pharmacy, supermarket or health food shop.**

**Some medicines may interfere with the way Lercadip works, or may be affected by Lercadip. These include:**

- ritonavir, ciclosporin, ketoconazole, itraconazole, erythromycin, fluoxetine, cimetidine (more than 800 mg daily).

You may need different amounts of your medicine, or you may need to take different medicines. Your doctor or pharmacist can tell you what to do if you are taking any of these medicines.

**Also ask your doctor or pharmacist what to do if you are taking, or are about to take the following medicines: phenytoin,**

**carbamazepine, rifampicin, amiodarone, quinidine, digoxin, simvastatin, metoprolol or propranolol.**

**If you have not told your doctor about any of the medicines listed above, tell them before you start taking Lercadip.**

### How to take Lercadip

#### *How much to take*

#### **Follow all directions given to you by your doctor carefully.**

These directions may differ from the information contained in this leaflet.

The usual dose is one 10 mg tablet taken once daily, but may be increased to 20 mg once daily.

If you do not understand the instructions on the box, ask your doctor or pharmacist for help.

#### *How to take it*

**Swallow Lercadip whole with a glass of water.**

#### *When to take it*

**Take Lercadip at about the same time each day, at least 15 minutes before a meal.**

Lercadip will have the best effect if it is taken at the same time each day at least 15 minutes before meals. This will also help you remember when to take the tablets.

#### *How long to take it*

Lercadip helps control your condition, but does not cure it. Therefore you must take Lercadip every day. Continue taking the tablets for as long as your doctor tells you to.

#### *If you forget to take it*

**If you forget to take a dose but remember within 12 hours from when the dose was due, take it straight away, then continue as normal the next day. Otherwise skip that day's dose but be sure to take the next day's dose when it is due.**

If you are not sure about what to do, talk to your doctor or pharmacist.

**Do not take a double dose to make up for the dose that you missed.**

If you have trouble remembering to take your tablets, ask your pharmacist for some hints.

If you have missed several doses, consult your doctor.

#### *If you take too much (overdose)*

**Immediately telephone your doctor or Poisons Information Centre (telephone 13 11 26 or New Zealand 0800 764 766), or go to Accident & Emergency at your nearest hospital, if you think that you or anyone else may have taken too much Lercadip.**

**Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention. Keep telephone numbers of these places/services handy.**

If you take too much Lercadip it may cause your blood pressure to become too low and you may feel your heart beats become irregular and faster. It may also lead to unconsciousness.

---

## While you are using Lercadip

---

### *Things you must do*

**Use Lercadip exactly as your doctor has prescribed.**

If you do not follow your doctor's instructions correctly, your blood pressure may not be controlled.

**Be sure to keep all of your doctor's appointments so that your progress can be checked.**

**If you become pregnant while you are taking Lercadip, tell your doctor.**

**If you are about to start taking any new medicine, tell your doctor and pharmacist that you are taking Lercadip.**

**Tell all other doctors, dentists and pharmacists who are treating you that you are taking Lercadip.**

**If you have an operation, tell the anaesthetist that you are taking this medicine.**

### *Things you must not do*

**Do not give Lercadip to anyone else, even if they have the same condition as you.**

**Do not use Lercadip to treat any other complaints unless your doctor tells you to.**

### *Things to be careful of*

**Be careful driving or operating machinery until you know how Lercadip affects you.**

Lercadip generally does not cause any problems with your ability to drive a car or operate machinery. However, as with other medicines used to treat high blood pressure, a few people may feel dizzy, light-headed or faint, especially when first taking Lercadip or when starting to take a different amount of medicine. Your doctor may also ask you to limit or stop your alcohol intake while taking medicines to control your blood pressure, such as Lercadip, as alcohol may increase these effects.

**If you feel light-headed, dizzy or faint when getting out of bed or standing up, get up slowly.**

**As with some medicines used to treat high blood pressure such as Lercadip, you should avoid drinking grapefruit juice as grapefruit juice may increase the effects of these medicines.**

---

## Side Effects

---

**Tell your doctor or pharmacist as soon as possible if you do not feel well while you are taking Lercadip.**

Lercadip helps most people with high blood pressure, but it may have unwanted side effects in a few people. All medicines can have side effects. Sometimes they are serious, most of the time they are not. You may need medical treatment if you get some of the side effects.

Possible side effects include:

- flushing
- swelling of the ankles, feet or lower legs
- palpitations
- headache
- dizziness
- gastrointestinal disturbances such as heartburn, nausea, epigastric pain or diarrhoea
- fatigue or sleepiness

These effects when they occur are usually mild. However, should you experience any of these or other undesirable effects not mentioned above, you should consult your doctor or pharmacist.

**Ask your doctor or pharmacist to answer any questions you may have.**

**Tell your doctor immediately or go to Accident & Emergency at your nearest hospital if you notice any of the following:**

- angina (chest pain or tightness)
- increased heart beat
- signs of allergy such as rash, itching or hives on the skin

These side effects are usually rare but may be serious and need urgent medical attention.

Do not be alarmed by this list of possible side effects. You may not experience any of them.

---

## After using Lercadip

---

### *Storage*

**Keep your tablets in the blister pack until it is time to take them.**

If you take the tablets out of the blister pack they may not keep well.

**Keep it in a cool dry place away from sunlight where the temperature stays below 30°C. Do not store it or any other medicine in the bathroom or near a sink.**

**Do not leave it in the car or on window sills.**

Heat and dampness can destroy some medicines.

**Keep it where children cannot reach it.**

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

### *Disposal*

**If your doctor tells you to stop taking the tablets or the tablets have passed their expiry date, ask your pharmacist what to do with any tablets that are left over.**

---

## Product description

---

### *What it looks like*

Lercadip is available in 10 mg or 20 mg tablets.

The 10mg film-coated tablets are yellow and round with a score line across the middle.

The 20mg film-coated tablets are round and pink.

Lercadip comes in packs of 28 tablets.

### *Ingredients*

#### **Active Ingredient**

Lercanidipine hydrochloride 10 mg or 20 mg per tablet.

#### **Excipient Ingredients**

Lactose monohydrate, microcrystalline cellulose, sodium starch glycollate, povidone, magnesium stearate, hypromellose, purified talc, titanium dioxide, macrogol 6000, Opadry OY-SR-6497 (yellow-10mg), Opadry O2-F2-5077 (red-20mg).

Lercadip does not contain gluten, sucrose, tartrazine or any other azo dyes.

---

## Sponsor

---

Lercadip 10 mg and 20 mg are supplied in Australia by:

Mylan Health Pty Ltd  
Level 1, 30 The Bond  
30-34 Hickson Road  
Millers Point, NSW 2000  
Australia

www.mylan.com.au

Phone: 1800 314 527

Australian registration numbers:

10 mg - AUST R 152709

20 mg - AUST R 152710

This leaflet was prepared on 20 March 2018