

Entecavir Mylan

contains the active ingredient (*Entecavir monohydrate*)

Consumer Medicine Information

What is in this leaflet

This leaflet answers some common questions about Entecavir Mylan.

It does not contain all the available information. It does not take the place of talking to your doctor or pharmacist.

All medicines have benefits and risks. Your doctor has weighed the risks of you taking Entecavir Mylan against the benefits expected for you.

If you have any concerns about taking this medicine, talk to your doctor or pharmacist.

Keep this leaflet with your medicine.

You may need to read it again.

What Entecavir Mylan is used for

This medicine is used to treat adults infected with hepatitis B virus.

This medicine belongs to a group of medicines called antiviral medicines.

Infection by hepatitis B virus can lead to damage to the liver. This medicine reduces the amount of virus in your body, and has been shown to improve the condition of the liver.

It is not known how safe Entecavir Mylan is when taken for long periods.

Ask your doctor if you have any questions about why this medicine has been prescribed for you.

Your doctor may have prescribed it for another reason.

This medicine is not addictive.

This medicine is available only with a doctor's prescription.

This medicine is not recommended for use in children under 16 years, as there have been no studies of its effects in children.

Before you take Entecavir Mylan

When you must not take it

Do not take Entecavir Mylan if you have an allergy to:

- any medicine containing entecavir monohydrate.
- any of the ingredients listed at the end of this leaflet.

Some of the symptoms of an allergic reaction may include:

- chills fever
- fast heart beat
- wheezing and coughing
- difficulty breathing
- dizziness
- flushing
- sweating and swelling of the face, lips, tongue or other parts of the body.

Do not take this medicine after the expiry date printed on the pack or if the packaging is torn or shows signs of tampering.

If it has expired or is damaged, return it to your pharmacist for disposal.

If you are not sure whether you should start taking this medicine, talk to your doctor.

Before you start to take it

It is important to remain under the care of your doctor during Entecavir Mylan therapy and after stopping Entecavir Mylan. You should report any new symptoms, medication or any other aspects affecting your health to your doctor. Your hepatitis B virus infection may get worse if you stop taking Entecavir Mylan. If your doctor advises you to stop Entecavir Mylan, they will monitor your health and perform regular blood tests to monitor your liver.

Tell your doctor if you have allergies to any other medicines, foods, preservatives or dyes.

Tell your doctor if you have or have had any of the following medical conditions:

- currently experience or have experienced any medical conditions especially any problems with your kidneys.
- have HIV and you are not currently on HIV treatment.
Entecavir Mylan is not recommended in patients who have both HIV and Hepatitis B and who are not currently receiving anti-HIV treatment. Entecavir Mylan may affect your HIV virus which could impact on future treatment options for HIV.
- are lactose intolerant. Entecavir Mylan contains lactose.

Tell your doctor if you are pregnant or plan to become pregnant or are breastfeeding.

Experience is limited with the use of Entecavir Mylan in pregnant women. Therefore, it should be used during pregnancy unless it is clearly needed. If there is an urgent need to consider Entecavir Mylan during pregnancy, your doctor will discuss with you the risks and benefits involved.

It is not known whether Entecavir Mylan passes into breast milk. Therefore to avoid possible side effects in the nursing infant, mothers should stop breastfeeding if they are taking Entecavir Mylan.

If you have not told your doctor about any of the above, tell him/her before you start taking Entecavir Mylan.

Taking other medicines

Tell your doctor or pharmacist if you are taking any other medicines, including any that you get without a prescription from a pharmacy, supermarket or health food shop.

Your doctor and pharmacist have more information on medicines to be careful with or avoid while taking this medicine.

How to take Entecavir Mylan

Follow all directions given to you by your doctor and pharmacist carefully.

They may differ from the information contained in this leaflet.

If you do not understand the instructions on the box/bottle, ask your doctor or pharmacist for help.

How much to take

The usual dose of Entecavir Mylan 0.5 mg or 1 mg is once a day.

If you have a medical problem with your kidneys your doctor may need to change how often you take your Entecavir Mylan tablets.

Your doctor will tell you what dose to take and how often you should take your Entecavir Mylan tablets.

How to take it

Swallow the tablets whole with a full glass of water.

The dose of Entecavir Mylan should be taken on an empty stomach.

When to take it

Take your medicine any time of day provided it is on an empty stomach, for example, at least 2 hours before food or 2 hours after food.

Talk to your doctor or pharmacist to work out when it is best for you to take your dose of Entecavir Mylan.

How long to take it

Continue taking your medicine for as long as your doctor tells you to.

This medicine helps to control your condition, but does not cure it. It is important to keep taking your medicine even if you feel well.

Entecavir Mylan is very important treatment that can improve the inflammation and scar tissue caused by the Hepatitis B virus in your liver and may reduce the change of developing cirrhosis, liver failure and liver cancer.

It is extremely important that you do not stop taking Entecavir Mylan without discussing it with your doctor. If Entecavir Mylan is suddenly stopped, the hepatitis B virus can become very active again and lead to sudden development of severe liver failure. There is high risk of dying if liver failure develops and liver transplantation may be necessary to save your life.

It is important to take Entecavir Mylan every day or as directed by your doctor, to not miss medicine doses, and to make sure you have enough supply until you next see your doctor.

Do not stop taking Entecavir Mylan or change the dose unless asked to do so by your doctor, even if you feel better, as it can be very dangerous.

If you forget to take it

If it is almost time for your next dose, skip the dose you missed and take your next dose when you are meant to.

Otherwise, take it as soon as you remember, and then go back to taking your medicine as you would normally.

Do not take a double dose to make up for the dose you missed.

This may increase the chance of you getting an unwanted side effect.

If you are not sure what to do, ask your doctor or pharmacist.

If you have trouble remembering to take your medicine, ask your pharmacist for some hints.

If you take too much (overdose)

Immediately telephone your doctor or the Poisons Information Centre (telephone 13 11 26) for advice, or go to Accident and Emergency at the nearest hospital, if you think that you or anyone else may have taken too much Entecavir Mylan. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

While you are taking Entecavir Mylan

Things you must do

If you are about to be started on any new medicine, remind your doctor and pharmacist that you are taking Entecavir Mylan.

Tell any other doctors, dentists and pharmacists who treat you that you are taking this medicine.

If you are going to have surgery, tell the surgeon or anaesthetist that you are taking this medicine.

It may affect other medicines used during surgery.

If you become pregnant while taking this medicine, tell your doctor immediately.

If you are about to have any blood tests, tell your doctor that you are taking this medicine. It may interfere with the results of some tests.

Keep all of your doctor's appointments so that your progress can be checked.

Things you must not do

Do not take Entecavir Mylan to treat any other complaints unless your doctor tells you to.

Do not give your medicine to anyone else, even if they have the same condition as you.

Do not stop taking your medicine or change the dosage without checking with your doctor. If you stop taking it suddenly, your condition may worsen.

Things to be careful of

Be careful driving or operating machinery until you know how Entecavir Mylan affects you.

This medicine may cause dizziness, in some people. It is not known if this was caused by Entecavir Mylan. If you have this symptom, do not drive, operate machinery or do anything else that could be dangerous.

If you feel light-headed, dizzy or faint when getting out of bed or standing up, get up slowly.

Standing up slowly, especially when you get up from bed or chairs, will help your body get used to the change in position and blood pressure. If this problem continues or gets worse, talk to your doctor.

Make sure that you visit your doctor regularly throughout your entire course of treatment with Entecavir Mylan.

When your treatment with Entecavir Mylan is stopped, your doctor will continue to monitor you and take blood tests for several months.

There is no evidence that Entecavir Mylan reduces the risk of infecting other with hepatitis B through sexual contact or body fluids (including blood contamination).

Therefore, it is important to take appropriate precautions to prevent others being infected with hepatitis B.

Talk to your doctor about safe sexual practices that protect your partner. Never share needles. Do not share personal items that can have blood or bodily fluids on them, like toothbrushes and razor blades. A vaccine is available to protect those at risk of becoming infected with hepatitis B.

Side effects

Tell your doctor or pharmacist as soon as possible if you do not feel well while you are taking Entecavir Mylan.

This medicine helps most people with hepatitis B infection, but it may have unwanted side effects in some people.

All medicines can have side effects. Sometimes they are serious, most of the time they are not. You may need medical attention if you get some of the side effects.

Do not be alarmed by the following list of side effects.

You may not experience any of them.

Ask your doctor or pharmacist to answer any questions you may have.

Tell your doctor or pharmacist if you notice any of the following and they worry you:

- diarrhoea
- indigestion
- tiredness
- headache.

The above list includes the more common side effects of your medicine.

Tell your doctor as soon as possible if you notice any of the following signs or symptoms of liver problems:

- skin and the white part of eyes turns yellow (jaundice)
- urine turns dark
- bowel movement (stool) turn light in colour
- loss of appetite
- nausea
- lower stomach pain
- lactic acidosis
- serious liver problems.

The above list includes serious side effects that may require medical attention.

Tell your doctor as soon as possible if you notice any of the following signs or symptoms of lactic acidosis:

- feeling very weak or tired
- unusual muscle pain
- trouble breathing
- stomach pain with nausea and vomiting
- feeling cold (especially in arms and legs)
- feeling dizzy or light-headed
- fast or irregular heartbeat.

The above list includes serious side effects that may require medical attention.

If any of the following happen, tell your doctor immediately or go to Accident and Emergency at your nearest hospital if you notice any of the following signs of a sudden life-threatening allergic reaction:

- chills
- fever
- fast heart beat
- wheezing or coughing
- difficulty breathing
- dizziness
- flushing
- sweating and swelling of the face, tongue or other parts of the body.

The above list includes very serious side effects. You may need urgent medical attention or hospitalisation.

Tell your doctor or pharmacist if you notice anything that is making you feel unwell.

Other side effects not listed above may also occur in some people.

After taking Entecavir Mylan

Storage

Keep your tablets until it is time to take them. If you take the tablets they may not keep well.

Keep your tablets in a cool dry place where the temperature stays below 25°C.

Do not store Entecavir Mylan or any other medicine in the bathroom or near a sink. Do not leave it on a window sill or in the car. Heat and dampness can destroy some medicines.

Keep it where children cannot reach it.

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

Disposal

If your doctor tells you to stop taking this medicine or the expiry date has passed, ask your pharmacist what to do with any medicine that is left over.

Product description

What it looks like

0.5 mg - A white, film-coated, round, biconvex, beveled edge tablet debossed with 'M' on one side of the tablet and 'EV1' on the other side. Blister pack of 30 tablets.

1 mg - A white, film-coated, round, biconvex, beveled edge tablet debossed with 'M' on one side of the tablet and 'EV2' on the other side. Blister pack of 30 tablets.

Ingredients

Entecavir Mylan 0.5 mg film coated tablet contains 0.5 mg of entecavir monohydrate as the active ingredient.

Entecavir Mylan 1 mg film coated tablet contains 1 mg of entecavir monohydrate as the active ingredient.

It also contains the following inactive ingredients:

- lactose
- microcrystalline cellulose
- crospovidone
- magnesium stearate
- Opadry White YS-1R-7003 (ARTG No. 1625).

Supplier

Entecavir Mylan is supplied in Australia by:

Alphapharm Pty Limited

(ABN 93 002 359 739)

Level 1, 30 The Bond

30-34 Hickson Road

Millers Point NSW 2000

Phone: (02) 9298 3999

www.alphapharm.com.au

Australian Registration Numbers:

Entecavir Mylan 0.5 mg: AUST R 220090

Entecavir Mylan 1 mg: AUST R 220091

This leaflet was prepared on 6 February 2015.

Entecavir Mylan_cmi\FEB15/00