

Patient Preparation Guidelines for SIBO Breath Testing

Procedures, Medications, and Supplements:

You must wait at least 1 month after colonoscopy or Barium studies, as well as antibiotic use before testing

One week prior to testing please stop the following

- Antacids/acid blocking medication
- Probiotics, such as Inner Health, or any probiotic containing product (yogurt, Yakult)
- Herbal antimicrobials

One day prior to testing, please stop

- Laxatives
- Digestive aids such as enzymes or hydrochloric acid

Test Kits

If you have ordered one test kit, please follow the instruction below. If you have 2 kits, resume the prep diet after completing the first test kit and use the second test kit the following morning.

For example,

Monday: prep diet followed by overnight fast


Tuesday AM: complete Lactulose test kit (3 hours) followed by resuming the prep diet and a 12 hour overnight fast

Wednesday AM: Glucose test kit (3 hours), then resume normal diet.

Prep Diet

Preparation begins 24 hours prior to testing; in 2 stages of 12 hours. If you suffer from constipation, please allow 48 hours prior to testing.

- First 12 hours consists of a restricted meal plan (see sample meal plan below), 36 hours if you are constipated
- Second 12 hours will be fasting – only water may be consumed;
- No smoking, including second-hand smoke, for at least 1 hour before or at any time during testing period;
- No sleeping or vigorous exercise for at least 1 hour before or at any time during testing period;



- Recent antibiotic therapy or runny diarrhoea may affect the results of the breath tests – please consult with your practitioner about these conditions prior to performing the test as testing may need to be rescheduled;
- Drinking water only during your breath test is allowed in moderation.

Sample Meal Plan

Only to be consumed during the first 12 hours, 48 hours if you are constipated.

- Baked or broiled chicken, fish or turkey (salt and pepper only)
- Plain steamed white rice (Basmati or Jasmine)
- Hard aged cheese (Parmesan, Pecorino) unless you are sensitive to dairy
- Eggs
- Clear chicken or beef broth
- Olive oil or coconut oil – 1 tablespoon only, for cooking

Avoid all other types of food during this 12 hour/48 hour period as this will ensure accurate test results.

Example

From: 8am til 8pm: 12 or 48 hours of restricted meal plan

Breakfast: Scrambled eggs with parmesan cheese

Mid morning snack: Chicken broth

Lunch: Chicken and white rice seasoned with salt and pepper

Mid afternoon snack: Chicken broth

Dinner: Fish and white rice seasoned with salt and pepper

Drinks: Water

From: 8pm til 8am: 12 hours fasting.

Water

Commence testing after 8am.

Note—you can set your own 12 hour overnight fasting



sibotest