

RESEARCH ARTICLE

Title:

**“JUST IN CASE”. USE OF LARGE BORE PERIPHERAL INTRAVENOUS CATHETERS
IN PARTURIENT WOMEN: A GLOBAL STUDY.**

Authors:

Prof Joan Webster^{1,2,3}

Dr Gillian Ray-Barruel^{2,3,4}

Prof Claire M Rickard^{1,2,3}

Ms Nicole Marsh^{1,2,3}

Mr Gabor Mihalía^{5,6}

Dr Evan Alexandrou^{2,7}

Affiliations

1. Royal Brisbane and Women’s Hospital, Herston, QLD 4006, Australia.
2. Alliance for Vascular Access Teaching and Research (AVATAR), Menzies Health Institute Queensland, Griffith University, Brisbane, Australia.
3. School of Nursing & Midwifery, Griffith University, Nathan, QLD, 4111 Australia
4. QEII Jubilee Hospital, Brisbane, Australia
5. School of Medicine, Griffith University, Nathan, QLD, 4111 Australia.
6. Centre for Applied Health Economics, Menzies Health Institute Queensland, Griffith University, Australia.
7. School of Nursing and Midwifery, Western Sydney University, Penrith, NSW. Australia
2751

Corresponding author

Name: Joan Webster

Address: Level 2, Bld 34, Royal Brisbane and Women's Hospital, Butterfield St,
Herston, QLD 4029, Australia

Email: joan.webster@health.qld.gov.au

Phone: +61 7 3646 8590

Word count 1,547

Key words

Blood Transfusion; Catheterisation, Peripheral; Postpartum Haemorrhage.

Disclosure statement

The authors wish to declare the OMG study received unrestricted investigator initiated research grants from Becton Dickinson (BD), CareFusion and 3M. B Braun provided funds for professional translation of data collection tools into several languages. All funds have been made payable to Griffith University or Western Sydney University and not to individual researchers.

Synopsis

Large bore, peripheral intravenous catheters are over-used for vascular access in perinatal women. These catheters may lead to vascular damage and increase phlebitis risk.

“JUST IN CASE”. USE OF LARGE BORE PERIPHERAL INTRAVENOUS CATHETERS IN PARTURIENT WOMEN: A GLOBAL STUDY.

ABSTRACT

Objective: The aim of the current study was to map a global picture of the practice of inserting large bore catheters in in parturient women.

Methods: We analysed a sub-set of data from a prospective, multi-centre, international prevalence study, the ‘One Million Global’ peripheral intravenous catheter study. Obstetric cohort was drawn from 163 maternity units in 35 countries (1477 women, 1577 catheters) and was collected between 1 June 2014 and 31 July 2015. Clinicians at each of the participating sites collected data using tools that had been previously validated. Results are reported as frequencies and proportions.

Results: Overall, 42% of all peripheral intravenous catheters were large bore and of all catheters placed in females in obstetric units; 438 (70%) of these were placed in the hand or wrist. The phlebitis rate was higher in the large bore group (12%) compared with those with smaller catheters (7%). Only 2% of women received blood products but it was unclear which catheter was used for this purpose.

Conclusion: Large bore peripheral intravenous catheters are over-used for vascular access in parturient women. They are painful and may cause vascular damage. If there is no indication, a catheter should not be placed at all.

INTRODUCTION

Approximately 830 women die from pregnancy or birth-related complications globally every day¹ with postpartum haemorrhage (PPH) being the major cause². Incidence of PPH is difficult to quantify due to different definitions and methods used to estimate blood loss, but recent large studies, where PPH is defined as a blood loss of ≥ 500 millilitres (ml), report a PPH rate of 1.2% to 9%³⁻⁷, with the highest rates in sub-Saharan Africa⁷. Variations in the inclusion and exclusion criteria of these studies may also contribute to disparity in the PPH rates.

Massive or severe PPH is defined variously as a blood loss $\geq 1,000$ ml^{4,7,8} or $\geq 1,500$ ml^{9,10}; bleeding requiring a blood transfusion⁵; or bleeding requiring ≥ 8 bags of red blood cells^{11,12}. The average rate for severe PPH is 1.2%, with a range of 0.02%–4.5%, depending on how PPH is defined^{4,5,7-12}. The most common cause of severe PPH, responsible for 27% of all maternal deaths², is uterine atony, which is the failure of the uterus to contract after the birth of a baby⁴. Consequently, to stimulate contractions, active management of the third stage of labour and use of oxytocin are widely recommended for PPH prophylaxis¹³.

Another common strategy used in Australia and New Zealand to prepare for negative sequelae from severe PPH is the prophylactic insertion of a large bore peripheral intravenous catheter (PIVC). In a recent Australian study, 88% of the 114 PIVCs placed in 95 perinatal women were 16 gauge or 18 gauge; 82% were placed in the hand or wrist; and 88% had extension tubing and a 3-way tap attached to the catheter¹⁴. Although these PIVCs are placed to facilitate rapid transfusion of blood products, the use of extension tubing and placement of the catheter in a small vein decreases the infusion flow rate by up to 76%^{15,16}. In addition, although most women in the study had a large bore catheter inserted, none were required for an urgent transfusion¹⁴.

Insertion of a large bore catheter, particularly in the hand or wrist is painful, around 4.5 on a 10-point scale¹⁷ and intravenous access guidelines recommend that the smallest gauge catheter should be selected to avoid vessel damage¹⁸, even when packed cells are transfused¹⁹. It is also likely, if a rapid transfusion was required, that a PIVC would be placed in the cubital fossa or another large vein. So, it remains unclear why the practice of placing large bore catheters in maternity patients persists. We also wondered if the practice was routine in other countries.

Aim:

To investigate the international prevalence of large bore PIVC use in parturient women

MATERIALS AND METHODS

Study design, patient population and setting

Our study is a sub-analysis from a prospective, multi-centre, international prevalence study, the One Million Global (OMG) PIVC study. Details of the study have been described elsewhere²⁰ but, briefly, hospitals were recruited using diverse strategies such as social media, network connections, conference presentations, and word of mouth. Adult and paediatric hospital in-patients with a PIVC *in situ* on the day of the study were eligible for inclusion; written or verbal consent was obtained from the patient or their next-of-kin at the time of data collection. Approval was granted by the Griffith University Human Research Ethics Committee (reference number NRS/34/13/HREC); ethics approval and/or managers approval was also required from each site prior to study commencement.

Data collection

Data was collected between 1 June 2014 and 31 July 2015. Clinicians at each of the participating sites collected data using tools that had been previously validated²¹. Items included catheter insertion characteristics, such as time, date and reason for insertion; catheter type and gauge; insertion site assessment, for signs of occlusion, infiltration, pain,

etc.; dressing type and integrity; and information about any infusates or intravenous medications. Phlebitis was defined as any pain, redness, swelling, purulent discharge, or palpable venous cord at the PIVC insertion site.

Statistical analysis

We defined large bore catheters as 14 to 18 gauge. Data management and analysis was undertaken using Stata 15 (StataCorp, College Station, Texas USA) statistical software. Results are reported as frequencies and proportions. Proportions were calculated using the number of non-missing observations in the denominator. Missing data were not imputed. In obstetric units, missing values of *gender* were recoded to *female*.

RESULTS

In the main study, a total of 40,620 PIVCs in 38,161 patients were assessed from 416 participating hospitals in 51 countries²¹. Data for the obstetric cohort was drawn from 163 maternity units in 35 countries. South Africa contributed the most sites (28) followed by Australia (25 sites). In total, 1,477 women accounted for 1,525 PIVCs in obstetric units. China studied the greatest number of PIVCs (245/1525) followed by Turkey (196/1525).

Large bore PIVCs comprised 624/1,493 (42%) of all PIVCs placed in females in obstetric units; 438 (70%) of these were placed in the hand or wrist. The phlebitis rate in obstetric units was 76/624 (12%) among those women with a large bore catheter in place compared with a rate of 65/869 (7%) in those with a smaller catheter. A total of 245/1506 (16%) PIVCs in females in obstetric units had no identified reason for use (no intravenous fluids or medications) on the day of the study. Of these idle catheters, 42 (17%) showed signs of phlebitis, and this rate was much higher than for PIVCs in active use (103/1,261; 8%). A total of 31 (2%) PIVCs were used for blood transfusion on the day of the study.

DISCUSSION

The aim of the current study was to investigate the international use of large bore PIVCs in parturient women. We found an overall prevalence of 42%, almost 2.5 times greater than the rate of 14-18G PIVCs among the 40,620 catheters in the overall cohort, which indicates an overuse in obstetric patients.

One of the recommendations for immediate treatment of severe PPH is to place an additional large gauge PIVC to facilitate the administration of red blood cells²². In this large, international cohort of parturient women, only 2.0% received a blood transfusion, although it is unclear how many of these were urgent and the size of the catheter used. The transfusion rate was similar to average rates from other sources^{4,5,7-12}, yet the fear of haemorrhage occurring exposes women to a 'just in case' approach and represents a pervasive and unnecessary invasive practice.

While placing a large bore catheter may seem an innocuous intervention, it carries many risks for the individual. The most important of these is vascular damage or scarring, which occurs when a large catheter is placed in a small vein, which may restrict future access opportunities and greatly increases the risk of thrombus formation². Additionally, phlebitis is more likely to be diagnosed in large bore catheters compared with smaller gauge catheters due to mechanical or chemical irritation of the endothelial layer^{18,23}. In our study a 67% higher rate of phlebitis was observed in those with large gauge catheters and, although rare, phlebitis has been associated with the much more serious condition, catheter-related blood stream infection²⁴.

In the current study, 70% of all PIVCs were placed in the hand or the wrist; apart from potential vascular damage and insertion pain, such veins are unsuitable for rapid blood infusion due to increased peripheral resistance from their smaller vessel diameter. Consequently, placement of a second or third 16–18-gauge PIVC in a large vein is

recommended in the case of a massive haemorrhage²²⁻²⁵. This recommendation implies, for the initial catheter, that a smaller gauge would have been adequate for the administration of fluids or medications in all women.

STRENGTHS AND LIMITATIONS

The major strength of the study was its size and the number of countries included; it represents the first attempt to quantify the international use of large bore PIVCs for vascular access in parturient women. The study was exposed to the usual limitations associated with prevalence studies. Data was collected on one day in each hospital, so results would have been more rigorous if a prospective cohort design had been used. For example, we only asked if the woman had received a blood transfusion on the day of data collection and if the reason for the PIVC was to transfuse blood. It is possible that the woman could have been transfused at an earlier time. However, irrespective of the study design, our PPH rate matched those from other studies, so we feel reasonably confident that they are representative of the populations from which they were drawn. We are also unsure if blood transfusions were for catastrophic bleeds, delivered under emergency conditions, or for other reasons. In the Australian study, all the transfusions were non-urgent, delivered over several hours and could have been delivered through a 20-gauge catheter because of a slower required rate for non-urgent transfusions¹⁴.

We did not have a denominator, that is, we do not know the number of women who were inpatients in maternity departments on the day of data collection; only the number who had a catheter *in situ*. Consequently, we cannot estimate the proportion of women in whom any catheter is placed.

CONCLUSION

Large bore PIVCs are over-used for vascular access in perinatal women. Considering the associated risks, careful attention should be given to inserting the right gauge catheter for

specific purposes and, if there is no indication, a catheter should not be placed in a vein at all.

REFERENCES

1. World Health Organization. Maternal Mortality Fact Sheet 2016 [updated 30/10/2017]. <http://www.who.int/mediacentre/factsheets/fs348/en/>. Accessed October 30 2017
2. Say L, Chou D, Gemmill A, et al. Global causes of maternal death: a WHO systematic analysis. *Lancet Glob Health*. 2014;2(6):e323-33.
3. Sheldon WR, Blum J, Vogel JP, et al. Postpartum haemorrhage management, risks, and maternal outcomes: findings from the World Health Organization Multicountry Survey on Maternal and Newborn Health. *BJOG*. 2014;121 Suppl 1:5-13.
4. Vendittelli F, Barasinski C, Pereira B, Lemery D, Group H. Incidence of immediate postpartum hemorrhages in French maternity units: a prospective observational study (HERA study). *BMC Pregnancy Childbirth*. 2016;16:242.
5. Kramer MS, Berg C, Abenhaim H, et al. Incidence, risk factors, and temporal trends in severe postpartum hemorrhage. *Am J Obstet Gynecol*. 2013;209(5):449 e1-7.
6. Bonnet MP, Basso O, Bouvier-Colle MH, et al. Postpartum haemorrhage in Canada and France: a population-based comparison. *PLoS One*. 2013;8(6):e66882.
7. Ononge S, Mirembe F, Wandabwa J, Campbell OM. Incidence and risk factors for postpartum hemorrhage in Uganda. *Reprod Health*. 2016;13:38.
8. Prick BW, Auf Altenstadt JF, Hukkelhoven CW, et al. Regional differences in severe postpartum hemorrhage: a nationwide comparative study of 1.6 million deliveries. *BMC Pregnancy Childbirth*. 2015;15:43.
9. Nyflot LT, Sandven I, Stray-Pedersen B, et al. Risk factors for severe postpartum hemorrhage: a case-control study. *BMC Pregnancy Childbirth*. 2017;17(1):17.
10. Dupont C, Occelli P, Deneux-Tharaux C, et al. Severe postpartum haemorrhage after vaginal delivery: a statistical process control chart to report seven years of continuous quality improvement. *Eur J Obstet Gynecol Reprod Biol*. 2014;178:169-75.

11. Green L, Knight M, Seeney FM, et al. The epidemiology and outcomes of women with postpartum haemorrhage requiring massive transfusion with eight or more units of red cells: a national cross-sectional study. *BJOG*. 2016;123(13):2164-70.
12. Ramler PI, van den Akker T, Henriquez D, Zwart JJ, van Roosmalen J. Incidence, management and outcome of women requiring massive transfusion after childbirth in the Netherlands: secondary analysis of a nationwide cohort study between 2004 and 2006. *BMC Pregnancy Childbirth*. 2017;17(1):197.
13. Dahlke JD, Mendez-Figueroa H, Maggio L, et al. Prevention and management of postpartum hemorrhage: a comparison of 4 national guidelines. *Am J Obstet Gynecol*. 2015;213(1):76 e1-10.
14. Webster J, Larsen E, Booker C, Laws J, Marsh N. Prophylactic insertion of large bore peripheral intravenous catheters in maternity patients for postpartum haemorrhage: A cohort study. *Aust N Z J Obstet Gynaecol*. 2017.
15. Levine DM, Garden AL, Truong HT, Bergemann J, Eames P. Influence of multi-lumen extensions on fluid flow through intravenous cannulae. *Anaesthesia*. 2013;68(12):1239-42.
16. Marks R, Boyd J, Ruiz N. Intravenous extension sets: when more is less. *J Clin Anesth*. 2013;25(4):348-50.
17. Tan PC, Mackeen A, Khong SY, Omar SZ, Azmi MA. Peripheral intravenous catheterisation in obstetric patients in the hand or forearm vein: a randomised trial. *Sci Rep*. 2016;6:23223.
18. Gorski L, Hadaway L, Hagle ME, Mc Goldrick M, Orr M, Doellman D. Infusion therapy standards of practice. *J Infus Nurs*. 2016;39(1S).
19. Stupnyckyj C, Smolarek S, Reeves C, McKeith J, Magnan M. Changing blood transfusion policy and practice. *Am J Nurs*. 2014;114(12):50-9.
20. Alexandrou E, Ray-Barruel G, Carr PJ, et al. Use of short peripheral intravenous catheters: characteristics, management, and outcomes worldwide. *J Hosp Med*. 2018;13(5).

21. Alexandrou E, Ray-Barruel G, Carr PJ, et al. International prevalence of the use of peripheral intravenous catheters. *J Hosp Med*. 2015;10(8):530-3.
22. Sentilhes L, Vayssiere C, Deneux-Tharaux C, et al. Postpartum hemorrhage: guidelines for clinical practice from the French College of Gynaecologists and Obstetricians (CNGOF): in collaboration with the French Society of Anesthesiology and Intensive Care (SFAR). *Eur J Obstet Gynecol Reprod Biol*. 2016;198:12-21.
23. Wallis MC, McGrail M, Webster J, et al. Risk factors for peripheral intravenous catheter failure: a multivariate analysis of data from a randomized controlled trial. *Infect Control Hosp Epidemiol*. 2014;35(1):63-8.
24. Mermel LA, Allon M, Bouza E, et al. Clinical practice guidelines for the diagnosis and management of intravascular catheter-related infection: 2009 Update by the Infectious Diseases Society of America. *Clin Infect Dis*. 2009;49(1):1-45.
25. National Institute for Health and Care Excellence. NICE Guideline 55 (2007) Intrapartum Care: care of healthy women and their babies during childbirth. Manchester: National Institute for Health and Clinical Excellence; 2007.